Roll Na	LHR-G1-12-19 (To be filled in b)
	(Academic Sessions 2015 - 2017 to 2017 - 2019)

filled in by the candidate)

PHYSICS

219-(INTER PART - II)

Time Allowed: 20 Minutes

Q.PAPER - II (Objective Type)

GROUP-I

Maximum Marks: 17

PAPER CODE = 8477

Note: Four possible answers A, B, C and D to each question are given. The choice which you think is correct, fill that circle in front of that question with Marker or Pen ink in the answer-book. Cutting or filling

two or more circles will re			
	30 - 10 00 E 10 - 10		
4			(D) hf
The sum of negative ar	nd positive peak values is	:	
(A) Average value	(B) rms value	(C) Peak value	(D) p-p value
		1 m ⁻¹ then the unit of	
		(C) ms	(D) ms ⁻¹
The common emitter co		rβ is given by:	
(A) $\frac{I_C}{I_C}$	(B) I_C	(C) $\frac{I_E}{I_E}$	(D) $\frac{I_B}{I_C}$
I_E	I_B	I_B	l_c
Resistance in choke is	:		The state of the s
(A) Large	(B) Very smail	(C) Zero	(D) Infinite
Sec/Ohm is equal to:			
(A) Farad	(B) Coulomb	(C) Joule	(D) Ampere
Number of neutrons in	²³⁵ ₉₂ U :		
(A) 92	(B) 235	(C) 143	(D) 327
Commutators are used			
		(C) A C motor	(D) A.C. rotator
			(1) 11.0.10401
The factor $\frac{1}{m}$ in Cor	npton equation has the di	mension of:	
	(R) Length	Mass	(D) Momentum
If a charged body is mo	oved against the electric f	ield, it will gain:	(1)
		ALL MARKET STATES STATES	rical notential energy
			ical potential energy
			(D) Noutrons
			(D) Neutrons
(A) Double	(B) Half	(C) Remain same	(D) One forth
		(0) 12 (.1/	(D) 0.95 -W
		(C) -13.6 eV	(D) - 0.85 eV
The gain of holl-invert		D	D
(A) 1+ 12	(B) $1 + \frac{\kappa_1}{R}$	(C) $\frac{-\kappa_2}{R}$	(D) $\frac{-R_1}{R_2}$
		Kı	Ka
N ₁	R ₂	4	
	nagnetic radiations having		e :
X-rays are the electron (A) $10^{-12}m$	R ₂	the wavelength in rang (C) $10^{-8}m$	
	nagnetic radiations having (B) $10^{-10} m$		e :
(A) $10^{-12}m$	nagnetic radiations having (B) 10 ⁻¹⁰ m transformer:	(C) 10 ⁻⁸ m	e :
(A) 10 ⁻¹² m To construct a step up	nagnetic radiations having (B) $10^{-10} m$ transformer: (B) $N_s < N_p$		e: (D) 10 ⁻⁶ m
(A) $10^{-12}m$ To construct a step up (A) $N_s > N_p$	nagnetic radiations having (B) $10^{-10} m$ transformer: (B) $N_s < N_p$	(C) $10^{-8}m$ (C) $N_s = N_p$	e: (D) 10 ⁻⁶ m
	The sum of negative ar (A) Average value The unit of \overline{E} is NC- (A) ms^{-2} The common emitter c (A) $\frac{I_C}{I_E}$ Resistance in choke is (A) Large Sec/Ohm is equal to: (A) Farad Number of neutrons in (A) 92 Commutators are used (A) D.C. generators The factor $\frac{h}{m_o c}$ in Cor (A) Pressure If a charged body is moderated as with the common emitter c (A) Pressure If a charged body is moderated by the common emitter c (A) Pressure If a charged body is moderated by the common emitter c (A) Pressure If a charged body is moderated by the common emitter c (A) Pressure If a charged body is moderated by the common emitter c (A) Pressure If a charged body is moderated by the common emitter c (A) P.E. (B) In p-type substances, the common emitter c (A) Double Energy of the 4 th orbit c (A) -2.51 eV The gain of non-invert	The sum of negative and positive peak values is (A) Average value (B) rms value The unit of \overline{E} is NC ⁻¹ and that of \overline{B} is NA ⁻¹ (A) ms^{-2} (B) $m^{-1}s^{-1}$ The common emitter current amplification factor (A) $\frac{I_C}{I_E}$ (B) $\frac{I_C}{I_B}$ Resistance in choke is: (A) Large (B) Very small Sec/Ohm is equal to: (A) Farad (B) Coulomb Number of neutrons in $\frac{235}{92}U$: (A) 92 (B) 235 Commutators are used in: (A) D.C. generators (B) A.C. generators The factor $\frac{h}{m_o c}$ in Compton equation has the difference of the charged body is moved against the electric form $\frac{h}{m_o c}$ (B) K.E. (C) Mechanic In p-type substances, the majority charge currier (A) Electrons (B) Protons When a wire of resistance R is cut into two equal (A) Double (B) Malf Energy of the 4 th orbit in hydrogen atom is: (A) -2.51 eV (B) -3.50 eV The gain of non-inverting amplifier is:	The sum of negative and positive peak values is: (A) Average value (B) rms value (C) Peak value The unit of \overline{E} is NC ⁻¹ and that of \overline{B} is NA ⁻¹ m ⁻¹ then the unit of (A) ms^{-2} (B) $m^{-1}s^{-1}$ (C) ms The common emitter current amplification factor β is given by: (A) $\frac{I_C}{I_E}$ (B) $\frac{I_C}{I_B}$ (C) $\frac{I_E}{I_B}$ Resistance in choke is: (A) Large (B) Very small (C) Zero Sec/Ohm is equal to: (A) Farad (B) Coulomb (C) Joule Number of neutrons in $\frac{235}{92}U$: (A) 92 (B) 235 (C) 143 Commutators are used in: (A) D.C. generators (B) A.C. generators (C) A.C. metor The factor $\frac{h}{m_o c}$ in Compton equation has the dimension of $\frac{1}{2}$: (A) Pressure (B) Length (C) Mass If a charged body is moved against the electric field it will gain: (A) P.E. (B) K.E (C) Mechanical energy (D) Electrons (D) Protons (C) Remain same Energy of the 4 th orbit in hydrogen atom is: (A) -2.51 eV (B) -3.50 eV (C) -13.6 eV The gain of non-finerting amplifier is:

To be filled in by the candidate) demic Sessions 2015 - 2017 to 2017 - 2019) PHYSICS 219-(INTER PART - II) Time Allowed: 2.40 hours PAPER - II (Essay Type) GROUP - I Maximum Marks: 68 SECTION-I 2. Write short answers to any EIGHT (8) questions : 16 (i) State Gauss's law and write its mathematical relation. (ii) Define electron volt and show that $1 eV = 1.6 \times 10^{-19} J$. (iii) Electric lines of force never cross. Why? (iv) Do electrons tend to go to region of high potential or of low potential? (v) State Lorentz force and write its formula. (vi) Write two uses of cathode ray oscilloscope. (vii) How can you use a magnetic field to separate isotopes of chemical element? (viii) Why the resistance of an ammeter should be very low? (ix) How the induced current can be increased? (x) What is motional emf and write its mathematical relation? (xi) Does the induced emf in a circuit depend on the resistance of the circuit? Explain III. (xii) Show that ε and $\frac{\Delta \phi}{\Delta t}$ have the same units. 16 (i) Define conventional current and solar cell. (ii) Define electrolysis and basic principle of electroplating. (iii) Why does the resistance of a conductor rise with temperature? (iv) Define peak value and peak to peak value of voltage or current. (v) A sinusoidal current has rms of 10A. What is the peak value?

3. Write short answers to any EIGHT (8) questions :

- (vi) What are superconductors?
- (vii) What is meant by para, diamagnetic substances?
- (viii) What is meant by strain energy
- (ix) Draw the truth table of XNOR gate.
- (x) Why ordinary silicon diodes do not emit light?
- (xi) Why is the base current in a transistor very small?
- (xii) Define intrinsic and extrinsic semi-conductor.

4. Write short answers to any SIX (6) questions :

(i) Will higher frequency light eject greater number of electrons than low frequency light?

- (ii) Photon A has twice the energy of photon B. What is the ratio of momentum of A to that of B?
- (iii) What is the energy of photon in a beam of infrared radiation of wavelength 1240 nm?
- (iv) What are the advantages of LASER over ordinary light?
- (v) Can the electron in ground state of hydrogen absorb a photon of energy 13.6 eV and greater than 13.6 eV?
- (vi) Define the isotopes of an element. Write down the isotopes of hydrogen.

(Turn Over)

12

LHR-12-G1 >

5

3

3

5

3

5

3


3

- 4. (vii) What is radioactive decay? Give an example.
 - (viii) What factor/make a fusion reaction difficult to achieve?
 - (ix) How can radioactivity help in the treatment of cancer?

SECTION - II

Note: Attempt any THREE questions.

- (a) State Gauss's law and apply it to find electric field intensity due to an infinite sheet of charge.
 - (b) A platinum wire has resistance of 10Ω at $0 \,^{\circ}$ C and 20Ω at $273 \,^{\circ}$ C. Find the value of temperature co-efficient of resistance.
- 6. (a) Define galvanometer. How it is converted into an ammeter and voltmeter?
 - (b) A pair of adjacent coils has a mutual inductance of 0.75 H. If the current in the primary changes from 0 to 10 A in 0.025 s, what is the average induced emf in the secondary? What is the change in flux in it, if the secondary has 500 turns?
- (a) Discuss the behaviour of an inductor in an A.C. circuit and write an expression for the inductive reactance.
 - (b) In circuit as shown in fig. there is negligible potential drop between B and E. If B is 100, calculate: (i) base current. (ii) collector current.
 - (iii) potential drop across R_c (iv) V_{CE}



- (a) Write down the postulates of special theory of relativity. Discuss the relation of time dilation, length contraction, mass variation and energy-mass relation with reference of this theory.
 - (b) A 1.0 m long copper wire is subjected to stretching force and its length increased by 20 cm Calculate the percent elongation which the wire undergoes.
- 9. (a) What are inner shell transitions? Describe the production of X-rays and their uses.
 - (b) How much energy is absorbed by a man of mass 80 kg who receives a lethal whole body equivalent dose of 400 rem in the form of low energy neutrons for which RBE factor is 10?

190-219-I-(Essay Type)-47000