Number: 4196 INTERMEDIATE P			Class) R	Roll No: MTN-2->		
	THEMATIC					
		ED: 30 Minutes e four choices for each object	OBJECT		XIMUM MAI	
	is correc	t, fill that bubble in front of t	ive type question hat question num	as A, B, C and D ber, on bubble s	heet. Use mark	nich you thin ker or ben to
	fill the b	ubbles. Cutting or filling two	or more bubbles	will result in zer	o mark in that	question.
.#	TI	QUESTIONS	A	В	С	D
1	$x^2 = -16y$	of directix of the parabola is:	y + 4 = 0	y - 4 = 0	x + 4 = 0	x - 4 = 0
2	The eccentric	eity of $\frac{y^2}{4} - x^2 = 1$ is:	$\frac{\sqrt{5}}{2}$	$\frac{2}{\sqrt{5}}$	$\frac{-2}{\sqrt{5}}$	2
3	$3\hat{i} \cdot (2\hat{j} \times \hat{k})$	= ?	0	2	3	6
4	$Cos \theta$ equal	to:	$\hat{a} \times \hat{b}$	$\hat{a}\cdot\hat{b}$	$ \hat{a} \times \hat{b} $	$\underline{a} \times \underline{b}$
5	The length of	the vector $2\hat{i} - 2\hat{j} - \hat{k}$ is:	3	4	5	2
6	The function and y is:	$x^2 + xy + y^2 = 2 \text{ of } x$	Constant	Even	Implicit	Explicit
7	If $f(x)=2x$	-8 , then $f^{-1}(x) = ?$	8 – 2 <i>x</i>	8+2x	$\frac{x-8}{2}$	x+8 2
8	$\frac{d}{dx}(3^x) = ?$		$\frac{3^{\prime}}{\ell n3}$	xln3	3× ℓn3	$3^{x} lnx$
)	If $y = \cos^{-1} \frac{1}{x}$	$\frac{x}{a}$, then $\frac{dy}{dx} = ?$	$\frac{-1}{\sqrt{a^2-x^2}}$	$\frac{-a}{\sqrt{x^2 + a^2}}$	$\frac{a}{\sqrt{x^2 - a^2}}$	$\frac{a}{\sqrt{a^2 - x^2}}$
0	$\frac{d}{dx}(\cos x) =$?	sinx	-sec x	sec x	$-\sin x$
1	If $y = \cos^{-1} \frac{y}{a}$	$\frac{x}{a}$, then $\cos y = ?$	The state of the s	$\frac{x}{a}$	$\frac{y}{a}$	sin y
2	$\int_{0}^{\pi} \sin x \ dx = ?$		$\cos \pi$	0	1	2
3	$\int \tan x \ dx =$?	$\ln \sec x + c$	$\ln \cos ec x + c$	$\ln \left \sin x \right + c$	$\ln \cot x + \epsilon$
4	$\int \frac{e^x}{e^x + 5} dx =$?	$(e^x+5)+c$	$\ln(e^x + 5) + c$	$e^{2x} + 5$	$e^{2x} + 7 + 6$
5	$\int -\cos ec^2x$	dx = ?	$\cos x + c$	$\tan x + c$	$\cos ec x + c$	$\cot x + c$
6		eclination of line ℓ , then $\frac{y_1}{\alpha} = r(say) \text{ is called:}$	Point-slope form	Normal form	Symmetric form	Two-point form
7	Equation of li third quadran	ne bisecting first and t is:	x = 0	<i>y</i> = 0	y = -x	y = x
8	The perpendic	cular distance of line = 0 from the origin is:	3	2	1	0
9	The graph of	$2x \ge 4$ lies in:	Upper Half	Lower Half	Left Half	Right Half
			Plane	Plane	Plane	Plane
0	Radius of circ	ele $x^2 + y^2 = 5$ is:	5	-5	$\sqrt{5}$	25

INTE	RMEDIATE PART-II (12th Class)	2024	(1 st -A)		Roll No:				
	HEMATICS PAPER-II GROUP-II	* * * * * * * * * * * * * * * * * * * *			DA A VINATURA DA A DIVIC. OO				
	E ALLOWED: 2.30 Hours SU E: Write same question number and its parts number	BJECTI		en in	MAXIMUM MARKS: 80				
NOI		TION-I							
2. At	tempt any eight parts.	N	1711-2	-2	8 × 2 = 16				
(i)	Define Implicit Function. (ii) Without finding the inverse, state the domain and range of f^{-1} $f(x) = \sqrt{x+2}$								
(iii)	Prove that $\lim_{x\to 0} (1+x)^{\frac{1}{x}} = e$	(iv)	Evaluate Lin	$\frac{\sin x}{x}$	<u>x⁰</u>				
(v)	Find by definition, derivative of $2x^2 + 1$ with respect to x		Differentiate	tiate with respect to 'x' $\frac{x^2+1}{x^2-3}$					
(vii)	Find $\frac{dy}{dx}$ if $y^2 - xy - x^2 + 4 = 0$	(viii)	Find $\frac{dy}{dx}$ if $x = y \sin y$ Find y_2 if $y = \ln\left(\frac{2x+3}{3x+2}\right)$						
(ix)	Find $f'(x)$ if $f(x) = x^3 e^{\frac{1}{x}}$, $x \neq 0$	(x)	Find y_2 if y	$y = \ln\left(\frac{2x+3}{3x+2}\right)$					
(xi)	By Maclaurin's series, show that $\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \dots$	(xii)	Determine in which interval 'f' is increasing or decreasing for domain mentioned $f(x)=4-x^2$, $x \in (-2,2)$						
3. At	tempt any eight parts.				8 × 2 = 16				
(i)	Find δy and dy in $y = x^2 - 1$ where x changes for	rom 3 to 3	3.02.						
(ii)	Evaluate the integral $\int \frac{1}{x^2 + 4x + 13} dx$	(iii)	Evaluate the integral $\int x \ln x dx$						
(iv)	Evaluate $\int_{-2}^{0} \frac{1}{(2x-1)^2} dx$	(v)	Find the area bounded by the curve $y = x^3 + 3x^2$ and the $x - $ axis.						
(vi)	Solve the differential equation $\sin y \cos ec x \frac{dy}{dx} = 1$	(vii)	Find the gener	Find the general solution of the equation $\frac{dy}{dx} - x = xy^2$					
(viii)	Show that the points $A(3, 1)$, $B(-2, -3)$ and $C(2, 2)$ are vertices of an isosceles triangle.								
(ix)	The xy - coordinate axes are rotated about the origin through the indicated angle and the new axes are OX and OY . Find the xy - coordinates of P with the given XY - coordinates $P(-5, 3)$; $\theta = 30^\circ$								
(v)			The same of the sa						
(x)	Write down an equation of the straight line passing through $(5, 1)$ (xi) Find the point of intersection of the lines and parallel to a line passing through the points $(0,-1)$, $(7,-15)$ (xi) Find the point of intersection of the lines $(5x+7y=35, 3x-7y=21)$								
(xii)	Find an equation of the line with x - intercept -3 and y - intercept 4.								
4. Att	tempt any nine parts.	10	de la constitución	1.2	9 2 = 18				
	Define Feasible Solution. (ii) Graph the inequality $x+2y<6$								
(iii)	Find the equation of the circle with centre at $(\sqrt{2}, -3\sqrt{3})$ and radius $2\sqrt{2}$.								
(iv)	Find focus and directix of the parabola $y^2 = -8(x-3)$								
(v)	Find length of tangent from the point $(-5, 10)$ to the ci	rcle $5x^2$	$+5y^2+14x-$	+123	y - 10 = 0				
(vi)	Find the centre and the foci of ellipse $9x^2 + y^2 = 18$ (vii) Write equation of hyperbola with foci (±5,0) and vertex (3,0).								
(viii)	Define Conic Section. (ix) Find the vector from the	ne point A	to the origin w	here 7	$\overrightarrow{AB} = 4\underline{i} - 2\underline{j}$ and B is the point $(-2, 5)$.				
(x)	If $ \alpha \underline{i} + (\alpha + 1)\underline{j} + 2\underline{k} = 3$. Find the value of α .								
(xi)	Show that the vectors $3\underline{i} - 2\underline{j} + \underline{k}$, $\underline{i} - 3\underline{j} + 5\underline{k}$ and $2\underline{i} + \underline{j} - 4\underline{k}$ form a right angle.								
(xii)	If $\underline{a} + \underline{b} + \underline{c} = 0$, then prove that $\underline{a} \times \underline{b} = \underline{b} \times \underline{c}$								
(xiii)	A force $F = 2\underline{i} + \underline{j} - 3\underline{k}$ acting at a point $A(1, -2, 1)$. Find the moment of \underline{F} about the point $B(2, 0, -2)$								
NOTE		II-NOI			3 × 10 = 30				
5.(a)				(b)					
	If $f(x) = \begin{cases} \frac{\sqrt{2x+5} - \sqrt{x+7}}{x-2}, & x \neq 2\\ k, & x = 2 \end{cases}$				Show that $\frac{dy}{dx} = \frac{y}{x}$ if $\frac{y}{x} = \tan^{-1} \frac{x}{y}$				
	Find value of 'k' so that 'f' is continuous at $x = 3$.				All and a second				
6.(a)	If $y = e^x \sin x$, show that $\frac{d^2y}{dx^2} - 2\frac{dy}{dx} + 2y = 0$			(b)	Evaluate $\int \sqrt{3-4x^2} dx$				
7.(a)	Evaluate $\int_{0}^{\sqrt{3}} \frac{x^3 + 9x + 1}{x^2 + 9} dx$ (b) Graph the feasible region of the following system of linear inequalities and find the corner points. $2x + 3y \le 18$, $2x + y \le 10$, $x + 4y \le 12$, $x \ge 0$, $y \ge 0$								
8.(a)	Find volume of the tetrahedron with vertices $A(2,1,8)$, $B(3,2,9)$, $C(2,1,4)$ and $D(3,3,10)$								
(b)	Write equations of two tangents from (2, 3) to the circle $x^2 + y^2 = 9$								
9.(a)	Show that the equation $9x^2 - 18x + 4y^2 + 8y - 23 = 0$ represents an ellipse. Find its elements.								
(b)	Find an equation of medians of the triangle whose vertices are $A(-3,2)$, $B(5,4)$ and $C(3,-8)$								

16-2024(1st-A)-12000 (MULTAN)