Paper Code 200 Number: 4181 INTERMEDIATE		23 (1 <sup>st</sup> -A) PART-II (12 <sup>th</sup> Class)		Roll No:			
STA	ATISTICS PA	PER-II	MTN-12-23				
TIN	ME ALLOWED	20 Minutes			MAXIMUM MARKS: 17		
Q.N	is correct, fill	r choices for each objective that bubble in front of tha es. Cutting or filling two or	t question num	ber, on bubl	ble sheet. Use mark	er or pen to	
S.#	OU	ESTIONS	A	В	С	D	
1		ibution, X lies between:	-∞ and 0	-∞ and +	$\infty$ 0 and $+\infty$	-1 and +1	
2	Standard normal random variable is denoted by:		Z	X	Y	ф	
3	If $X \sim N(100, 25)$ , then median is:		25	5	100	. 0	
4	Sample size is denoted by:		N	m	N(N-1)	n	
5	Any measure of the sample is called:		Parameter	Statistic	Constant	Sampling	
6	The random digits from 0 to 9 have probability:		1/10	$\frac{2}{10}$	10	1 100	
7	The statistical inference can be divided into approaches:		Four	Three	Two	Five	
8	Level of confidence is denoted by:		· «5	β	1-β	1-α	
9	P (Rejecting $H_0/H_0$ is true) is equal to:		$1-\alpha$	α	1+α	β	
10	Dependent variable is also called:		Predictor	Regresso	or Regressand	Fixed	
11	The sum of the re	The sum of the residuals is:		One	Two	Three	
12	When two variables are uncorrelated the $r$ is:		-1	+1	2	0	
13	For a contingency table $d.f. = 12$ and $r = 4$ , the $c$ is:		4	5	2	3	

40(Obj)(☆)-2023(1<sup>st</sup>-A)-2500 (MULTAN)

-1 and +1

Histogram

Cyclical

variations

RAM

-1 and 0

Bar diagram

Seasonal

variations

ROM

0 and 1

Historigram

Secular

trend

ALU

Rank correlation coefficient lies

The graph of time series is called:

The brain of the computer is known as:

A rise in prices before eid is

14

15

16

17

between:

an example of:

-2 and 1

Curve

Irregular

variations

CPU

INTERMEDIATE PART-II (12th Class)	2023 (1 <sup>st</sup> -A)	Roll No:
STATISTICS PAPER-II	MTN-12-2	?
TIME ALLOWED: 2.40 Hours		MAXIMUM MARKS: 68

		ECTION	I-I
2. A	ttempt any eight parts.		8 × 2 = 16
(i)	Define normal probability distribution.	(ii)	Express the term standard normal variate.
(iii)	Enlist two properties of normal distribution.	(iv)	In a normal distribution, $\mu = 163$ , $Q_3 - 171.094$ . Compute standard deviation of the given distribution.
(v)	In a normal distribution, $\mu = 24$ , $\sigma = 4$ . Calculate fourth moment about mean.	(vi)	Describe the term statistical inference.
(vii)	Define an estimator.	(viii)	Elaborate type – II error.
(ix)	Define level of significance.	(x)	Given $n = 16$ , $s = 0.75$ , $\overline{X} = 10.5$ , $\mu_0 = 10$ . Compute the test statistic (t - test).
(xi)	Describe a monitor.	(xii)	What is meant by byte?
3. At	tempt any eight parts.	J (All)	8 × 2 = 16
(i)	Define sample.	(ii)	What is standard error?
(iii)	Define the term bias.	(iv)	What is non-sampling error?
(v)	Given $n = 9$ , $\mu_{\overline{X}} = 4$ , $\sigma_{\overline{X}} = 2.5$ find $\mu$ and $\sigma$ .	(vi)	If $n = 40$ , $\pi = 0.7$ then find $\mu_P$ and $\sigma_P$ .
(vii)	What is meant by regression?	(viii)	Define dependent variable.
(ix)	Define the term correlation.	(x)	Interpret the meaning when $r = -1$ .
(xi)	Given that $\overline{x} = 1$ , $\overline{y} = 8$ , $b = 2$ find $y - $ intercept.	(xii)	If $\hat{y} = 11.8 + 2x$ and $\hat{x} = -5.5 + 0.5y$ then find $r$ .
4. At	tempt any six parts.		6 × 2 = 12
(i)	Define a contingency table.		
(ii)	Given $(AB) = 95$ , $(A\beta) = 55$ , $(\alpha\beta) = 85$	and $(\alpha\beta)$	) = 45. Find the coefficient of association.
(iii)	Given $f_0 = 7, 8, 15, 20$ and $f_e = 11.88, 12.8$		
(iv)	Define a time series.		
(v)	Explain the term "Noise" in time series.	7	
(vi)	Explain what is meant by seasonal variations?		
(vii)	Given $(Y - \hat{Y}) = 0.5, -0.5, 1, -1, 0.5, -0.5$	5. Find s	sum of squares of residuals.
(viii)	Given $\hat{Y} = 10 + 3X$ find the trend values for		
(ix)	What do you mean by Historigram?		

## SECTION-II

<u>SECTION-II</u>					
E: Attempt any three questions. $3 \times 8 = 24$					
<ul> <li>If the diameters of ball bearings are normally distributed with mean 0.6140 inches and standard deviation 0.002 inches. Determine the percentage of ball bearings with diameters.</li> <li>(i) less than 0.608 inches</li> <li>(ii) greater than 0.617 inches</li> <li>Scores on a national education achievement test are normally distributed with μ = 500 and σ = 100</li> <li>(i) What is the 95<sup>th</sup> percentile of this distribution?</li> <li>(ii) What are the lower and upper quartiles of this distribution?</li> </ul>					
If $N_1 = 400$ , $N_2 = 200$ , $n_1 = 100$ , $n_2 = 110$ , $\mu_1 = 500$ , $\mu_2 = 800$ , $\sigma_1 = 10$ , $\sigma_2 = 10$ obtain mean and standard error of sampling distribution of $\overline{X}_1 - \overline{X}_2$ . If sampling is done W.O.R.					
If $\bar{x} = 100$ , $s = 8$ and $n = 64$ . Construct a 99% confidence interval for population mean $(\mu)$ .					
A random sample of 25 values gives the average 83. Can this sample regarded as drawn from the normal population with mean 80 and $\sigma = 7$ with $\alpha = 0.05$					
Calculate correlation coefficient and interpret it between marks and study hours.  Marks 10 15 9 21 7 Study Hours 2 3 1 4 1					
Fit a regression line to data given in part(a) to predict marks. Estimate marks when study hours are 5.					
Given the following data. Find whether $A$ and $B$ are independent or associated. $n = 150$ ; $(A) = 30$ ; $(B) = 60$ ; $(AB) = 12$					
If the linear trend in the data for the years 1960 to 1965. Both inclusive with origin at the middle of 1962 and 1963 is $\hat{y} = 1306.667 + 73.428x$ , the unit of x being one year, then determine the trend line with origin at 1960 and hence determine the trend values.					