

MATHEMATICS

OBJECTIVE

TIME: 30 MINUTES

GROUP : FIRST

DGK-11-1-23

MARKS: 20

NOTE : You have four choices for each objective type question as A, B, C and D. The choice which you think is correct, fill that circle in front of that question number. Use marker or pen to fill the circles. Cutting or filling two or more circles will result in zero mark in that question.

QUESTION NO. 1

- 1 A.M between $\frac{1}{2}$ and $\frac{1}{4}$ is
(A) $-\frac{1}{8}$ (B) $\frac{1}{8}$ (C) $\frac{3}{4}$ (D) $\frac{3}{8}$
- 2 If $r = n$ then nC_r is equal to
(A) 0 (B) 1 (C) $n!$ (D) $(n-1)!$
- 3 For mutually exclusive events A and B
(A) $A \cup B = \emptyset$ (B) $A - B = \emptyset$ (C) $A \cap B = \emptyset$ (D) $A \cup B = A \cap B$
- 4 The in-equality $n^2 > n + 3$ is valid if
(A) $n \geq 2$ (B) $n \geq 0$ (C) $n \geq 1$ (D) $n \geq 3$
- 5 Sum of even coefficient in expansion of $(a+b)^4$ is
(A) 18 (B) 10 (C) 12 (D) 16
- 6 The angle $\frac{\pi}{12}$ in degree measure is
(A) 30° (B) 20° (C) 45° (D) 15°
- 7 $\sin 390^\circ$ is equal to
(A) $\cos 30^\circ$ (B) Zero (C) $\sin 30^\circ$ (D) $\sin 60^\circ$
- 8 Smallest positive number 'p' for which $f(x+p) = f(x)$ is called
(A) Domain (B) Range (C) Co-domain (D) Period
- 9 Radius of e-circle opposite to vertex B of triangle ABC is
(A) $\frac{\Delta}{s-a}$ (B) $\frac{\Delta}{s-b}$ (C) $\frac{\Delta}{s-c}$ (D) $\frac{\Delta}{s}$
- 10 In an equilateral Triangle ABC $r_1 : r_2 : r_3$ is equal to
(A) 1 : 2 : 3 (B) 1 : 3 : 3 (C) 3 : 3 : 3 (D) 2 : 3 : 3
- 11 $\cos^{-1}(-x) = ?$
(A) $\pi - \cos^{-1}x$ (B) $\cos^{-1}x$ (C) $\pi + \cos^{-1}x$ (D) $\sin^{-1}x$
- 12 Solution of $\tan 2x = 1, x \in [0, 2\pi]$ is
(A) $\left\{\frac{\pi}{8}, \frac{5\pi}{8}\right\}$ (B) $\left\{\frac{\pi}{4}, \frac{3\pi}{4}\right\}$ (C) $\left\{\frac{\pi}{4}, \frac{5\pi}{4}\right\}$ (D) $\left\{\frac{\pi}{6}, \frac{5\pi}{6}\right\}$
- 13 $(-i)^{19}$ is equal to
(A) $-i$ (B) 1 (C) -1 (D) i
- 14 A function $f: A \rightarrow B$ is surjective if
(A) Range of $f = A$ (B) Range of $f = B$ (C) Range of $f \neq B$ (D) Both A and B
- 15 A matrix $m \times 1$ is called
(A) Scalar Matrix (B) Row Matrix (C) Column Matrix (D) Null Matrix
- 16 If 'A' is a square Matrix of order 2×2 then $|KA|$ is equal to
(A) $2K|A|$ (B) $K^3|A|$ (C) $K|A|$ (D) $K^2|A|$
- 17 If one solution of equation $x^2 - ax + 2 = 0$ is $x = 1$ the 'a' is equal to
(A) 0 (B) -7 (C) 7 (D) 3
- 18 A quadratic equation $ax^2 + bx + c = 0$ becomes linear if
(A) $a = 0, b \neq 0$ (B) $a \neq 0$ (C) $b = 0$ (D) $b \neq 0$
- 19 $\frac{A}{x-1} + \frac{B}{x+1}$ are partial fractions of
(A) $\frac{1}{x^3-1}$ (B) $\frac{1}{x^2-1}$ (C) $\frac{1}{x^2+1}$ (D) $\frac{1}{x^3+1}$
- 20 $\sum_{k=1}^n k$ is equal to
(A) $\frac{n^2(n+1)^2}{4}$ (B) $\frac{n(n+1)(2n+1)}{6}$ (C) $\frac{n(n+1)}{2}$ (D) $\frac{n^2(n+1)}{4}$

DOK-11-1-23

QUESTION NO. 2 Write short answers of any Eight (8) parts of the following

16

i	State commutative law of addition and associative law of multiplication of real numbers.
ii	Separate into real and imaginary parts $\frac{4}{1+i}$
iii	Write the set $\{x/x \in \mathbb{R} \wedge x \neq x\}$ in the descriptive and tabular form
iv	Write converse and inverse of the conditional $\sim p \rightarrow q$
v	Show that the statement $(p \wedge q) \rightarrow p$ is tautology.
vi	If $A = \begin{bmatrix} 1 & -1 \\ a & b \end{bmatrix}$ and $A^2 = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$, find the values of "a" and "b"
vii	If $A = \begin{bmatrix} 1 & 2 & -3 \\ 0 & -2 & 0 \\ -2 & -2 & 1 \end{bmatrix}$, then find A_{32}
viii	If the matrices A and B are symmetric and $AB = BA$, show that AB is symmetric
ix	Define reciprocal equation.
x	Evaluate $(1 + \omega - \omega^2)^8$
xi	Prove that sum of four 4th roots of unity is zero.
xii	Use remainder theorem to find the remainder when $x^2 + 3x + 7$ is divided by $x + 1$

QUESTION NO. 3 Write short answers of any Eight (8) parts of the following

16

i	What are partial fractions ?
ii	Find the 13th term of the sequence $x, 1, 2-x, 3-2x, \dots$
iii	Find three A.Ms between 3 and 11.
iv	The sum of S_9 and S_7 is 203 and $S_9 - S_7 = 49$, S_7 and S_9 being the sums of the first 7 and 9 terms of an A.P respectively. Determine the series.
v	If $\frac{1}{a}, \frac{1}{b}$ and $\frac{1}{c}$ are in G.P show that the common ratio is $\pm \sqrt{\frac{a}{c}}$
vi	Find the Geometric means between 4 and 16.
vii	Find the value of n when ${}^n P_4 : {}^{n-1} P_3 = 9 : 1$
viii	In how many ways can 4 keys be arranged on a circular key ring?
ix	A natural number is chosen out of first fifty natural numbers. What is the probability that the chosen number is a multiple of 3 or of 5?
x	Prove the formula for $n = -1, 0, 1, 2, \dots$ $3+5+7+\dots+(2n+5) = (n+2)(n+4)$
xi	Expand $(a - \sqrt{2}x)^4$
xii	Expand the following up to 4 terms $(2 - 3x)^{-2}$

QUESTION NO. 4 Write short answers of any Nine (9) parts of the following

18

i	Show that the area of a sector of a circular region of radius r is $\frac{1}{2} r^2 \theta$, where θ is the circular measure of central angle of the sector.
ii	If $\tan \theta = \frac{1}{\sqrt{7}}$ and the terminal arm of the angle is not in III quad, find the value of $\frac{\operatorname{Cosec}^2 \theta - \sec^2 \theta}{\operatorname{Cosec}^2 \theta + \sec^2 \theta}$
iii	Prove the identity $(\sec \theta + \tan \theta)(\sec \theta - \tan \theta) = 1$
iv	Prove that $\cos 330^\circ \sin 600^\circ + \cos 120^\circ \sin 150^\circ = -1$
v	Prove $\frac{\sin(\alpha+\beta)+\sin(\alpha-\beta)}{\cos(\alpha+\beta)+\cos(\alpha-\beta)} = \tan \alpha$
vi	Prove the identity $\cot \alpha - \tan \alpha = 2 \cot 2\alpha$
vii	Find the period of $\sec 9x$
viii	Find the area of ΔABC , given three sides $a=18, b=24, c=30$
ix	Show that the $r_3 = s \tan \frac{Y}{2}$
x	Prove that $\tan \frac{\alpha}{2} = \sqrt{\frac{(s-b)(s-c)}{s(s-a)}}$
xi	Without using Calculator show that $\cos^{-1} \frac{4}{5} = \cot^{-1} \frac{4}{3}$
xii	Solve $\sec^2 \theta = \frac{4}{3}$, $\theta \in [0, 2\pi]$
xiii	Find the value of θ $2 \sin^2 \theta - \sin \theta = 0$, $\theta \in [0, 2\pi]$

SECTION-II

DGK-11-1-23

Note: Attempt any Three questions from this section

10 x 3 = 30

Q. 5-(A)	Show that $\begin{vmatrix} x & 1 & 1 & 1 \\ 1 & x & 1 & 1 \\ 1 & 1 & x & 1 \\ 1 & 1 & 1 & x \end{vmatrix} = (x+3)(x-1)^3$
(B)	Solve the equation $\left(x + \frac{1}{x}\right)^2 - 3\left(x + \frac{1}{x}\right) - 4 = 0$
Q. 6 -(A)	Resolve $\frac{1}{(x-1)^2(x+1)}$ into partial fraction
(B)	Prove that ${}^{n-1}C_r + {}^{n-1}C_{r-1} = {}^nC_r$
Q. 7-(A)	If $y = \frac{x}{2} + \frac{x^2}{4} + \frac{x^3}{8} + \dots$ and if $0 < x < 2$ then prove that $x = \frac{2y}{1+y}$
(B)	Identify the series : $1 + \frac{1}{3} + \frac{1.3}{3.6} + \frac{1.3.5}{3.6.9} + \dots$ as a binomial expansion and find its sum.
Q. 8 -(A)	If $\cot \theta = \frac{5}{2}$ and the terminal arm of the angle is in 1st quadrant. Find the values of $\frac{3 \sin \theta + 4 \cos \theta}{\cos \theta - \sin \theta}$
(B)	Prove that $\frac{2 \sin \theta \sin 2 \theta}{\cos \theta + \cos 3 \theta} = \tan 2 \theta \tan \theta$
Q. 9 -(A)	Prove that $r_1 + r_2 + r_3 - r = 4R$
(B)	Prove that $\cos^{-1} \frac{63}{65} + 2 \tan^{-1} \frac{1}{5} = \sin^{-1} \frac{3}{5}$