

**MATHEMATICS PAPER-I**  
**GROUP-II**

MTN-42-21

**OBJECTIVE**

TIME ALLOWED: 30 Minutes

MAXIMUM MARKS: 20

**Note:** You have four choices for each objective type question as A, B, C and D. The choice which you think is correct, fill that bubble in front of that question number, on bubble sheet. Use marker or pen to fill the bubbles. Cutting or filling two or more bubbles will result in zero mark in that question. No credit will be awarded in case BUBBLES are not filled. Do not solve question on this sheet of OBJECTIVE PAPER.

**Q.No.1**

- (1) The A.M between  $\sqrt{2}$ ,  $3\sqrt{2}$  is: (A)  $\sqrt{6}$  (B)  $-2\sqrt{2}$  (C)  $2\sqrt{2}$  (D)  $-\sqrt{6}$
- (2) Common ratio of G.P  $\frac{1}{a}, \frac{1}{b}, \frac{1}{c}$  is:  
 (A)  $\pm \sqrt{\frac{a}{c}}$  (B)  $\pm \sqrt{\frac{c}{a}}$  (C)  $\pm \sqrt{\frac{b}{c}}$  (D)  $\pm \sqrt{\frac{c}{b}}$
- (3) H.M between 3 and 7 is: (A) 5 (B)  $\sqrt{21}$  (C)  $\frac{21}{5}$  (D)  $\frac{5}{21}$
- (4) If A and B are two independent events, then  $P(A \cap B) =$   
 (A)  $P(A) + P(B)$  (B)  $P(A) - P(B)$  (C)  $P(A \cup B)$  (D)  $P(A) \cdot P(B)$
- (5) The number of terms in the expansion of  $(a+x)^n$  are:  
 (A) n (B) n+1 (C) n-1 (D) 2n
- (6) The value of  $\tan \theta$  for  $\theta = 30^\circ$  is: (A)  $\sqrt{3}$  (B)  $\frac{1}{\sqrt{3}}$  (C)  $\frac{2}{\sqrt{3}}$  (D)  $\frac{\sqrt{3}}{2}$
- (7)  $\frac{5\pi}{6}$  radian = (A)  $150^\circ$  (B)  $130^\circ$  (C)  $120^\circ$  (D)  $60^\circ$
- (8) If  $\sin \alpha = \frac{4}{5}$ ,  $0 < \alpha < \frac{\pi}{2}$ , then  $\cos \alpha =$  (A)  $\frac{2}{5}$  (B)  $\frac{1}{5}$  (C)  $\frac{4}{5}$  (D)  $\frac{3}{5}$
- (9)  $\pi$  is the period of: (A)  $\sec \theta$  (B)  $\operatorname{cosec} \theta$  (C)  $\cot \theta$  (D)  $\sin 3\theta$
- (10) In any triangle ABC, with usual notation  $\sqrt{\frac{s(s-c)}{ab}} =$   
 (A)  $\cos \frac{\gamma}{2}$  (B)  $\cos \frac{\alpha}{2}$  (C)  $\cos \frac{\beta}{2}$  (D)  $\sin \frac{\alpha}{2}$
- (11) Radius of e-circle opposite to vertex 'A' of  $\Delta ABC$  is:  
 (A)  $\frac{\Delta}{s-a}$  (B)  $\frac{\Delta}{s-c}$  (C)  $\frac{\Delta}{s}$  (D)  $\frac{\Delta}{s-b}$
- (12)  $2 \tan^{-1}(A) =$   
 (A)  $\tan^{-1}\left(\frac{2A}{1-A^2}\right)$  (B)  $\tan^{-1}\left(\frac{A}{1+A^2}\right)$  (C)  $\tan^{-1}\left(\frac{2A}{1-A^2}\right)$  (D)  $\tan^{-1}\left(\frac{2A}{1+A^2}\right)$
- (13) Reference angle of  $\sin x = \frac{1}{2}$  is: (A)  $\frac{\pi}{3}$  (B)  $\frac{\pi}{6}$  (C)  $\frac{\pi}{4}$  (D)  $\frac{\pi}{2}$
- (14) Every non terminating, non recurring decimal represents:  
 (A) Rational number (B) Irrational number (C) Natural number (D) Whole number
- (15) If A and B are any subsets of U, then  $A - B =$   
 (A)  $A \cup B^c$  (B)  $(A \cup B)^c$  (C)  $(A \cap B)^c$  (D)  $A \cap B^c$
- (16) A square matrix  $A = [a_{ij}]$  is called upper triangular matrix if:  
 (A)  $a_{ij} = 0$  for  $i < j$  (B)  $a_{ij} = 0$  for  $i > j$  (C)  $a_{ij} \neq 0$  for  $i > j$  (D)  $a_{ij} = k$  for  $i < j$
- (17) The trivial solution of system of homogeneous linear equation in three variables is:  
 (A) (0, 0, 1) (B) (0, 1, 0) (C) (0, 0, 0) (D) (0, -1, 0)
- (18) If  $\alpha, \beta$  are the roots of  $x^2 - px - p - c = 0$ , then  $\alpha\beta =$   
 (A)  $-p-c$  (B)  $p+c$  (C)  $p-c$  (D)  $-p+c$
- (19) Sum of all the four fourth roots of unity is: (A) 1 (B) 0 (C) -1 (D) 2
- (20) Partial fraction of  $\frac{x^2+1}{(x+1)(x-1)}$  will be of the form:  
 (A)  $\frac{A}{x-1} + \frac{B}{x+1}$  (B)  $\frac{A}{x+1} + \frac{Bx+C}{x-1}$  (C)  $\frac{Ax+B}{x^2-1}$  (D)  $1 + \frac{A}{x+1} + \frac{B}{x-1}$

INTERMEDIATE PART-I (11<sup>th</sup> CLASS)

MATHEMATICS PAPER-I

MTN-42-21

TIME ALLOWED: 2.30 Hours

GROUP-II

SUBJECTIVE

MAXIMUM MARKS: 80

**NOTE:** Write same question number and its part number on answer book, as given in the question paper.

SECTION-I

2. Attempt any eight parts.

8 × 2 = 16

- (i) Prove the following rule of addition  $\frac{a}{c} + \frac{b}{c} = \frac{a+b}{c}$
- (ii) Find the multiplicative inverse of  $(-4, 7)$
- (iii) If  $Z_1$  and  $Z_2$  are the complex numbers then prove that  $|Z_1 \cdot Z_2| = |Z_1| \cdot |Z_2|$
- (iv) Write the set  $\{x \mid x \in N \wedge x \leq 10\}$  into (i) Descriptive form (ii) Tabular form
- (v) Determine that  $p \rightarrow (p \vee q)$  is a tautology or not.
- (vi) Find the domain and range of the relation  $\{(x, y) \mid x + y > 5\}$  if  $A = \{1, 2, 3, 4\}$
- (vii) Find  $x$  and  $y$  if  $\begin{bmatrix} x+3 & 1 \\ -3 & 3y-4 \end{bmatrix} = \begin{bmatrix} 2 & 1 \\ -3 & 2 \end{bmatrix}$
- (viii) If  $A$  and  $B$  are two square matrices of same order, then explain why in general  $(A+B)(A-B) \neq A^2 - B^2$

(ix) Without expansion, show that  $\begin{vmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{vmatrix} = 0$

(x) Find four fourth roots of unity.

(xi) Find the number, if sum of a positive number and its reciprocal is  $\frac{26}{5}$

(xii) Discuss the nature of the roots of the equation  $2x^2 + 5x - 1 = 0$

3. Attempt any eight parts.

8 × 2 = 16

- (i) Resolve into partial fractions  $\frac{1}{(x-1)(2x-1)(3x-1)}$
- (ii) Resolve into partial fractions, without finding the constants  $\frac{x^2+15}{(x^2+2x+5)(x-1)}$
- (iii) If  $\frac{1}{a}$ ,  $\frac{1}{b}$  and  $\frac{1}{c}$  are in A.P, show that  $b = \frac{2ac}{a+c}$
- (iv) How many terms of the series  $-7 + (-5) + (-3) + \dots$ , amount to 65?
- (v) Find geometric means between 2 and 16.
- (vi) If  $y = \frac{x}{2} + \frac{1}{4}x^2 + \frac{1}{8}x^3 + \dots$  and if  $0 < x < 2$ , prove that  $x = \frac{2y}{1+y}$
- (vii) Prove that  ${}^nP_r = n \cdot {}^{n-1}P_{r-1}$
- (viii) How many arrangements of the letters of word, taken all together, can be made "PAKISTAN"?
- (ix) What is the probability that a slip of numbers divisible by 4 are picked from the slips bearing numbers 1, 2, 3, ..., 10?
- (x) Show that the inequality  $4^n > 3^n + 4$  is true for integral values of  $n = 2, 3$
- (xi) Expand upto three terms  $(4-3x)^{\frac{1}{2}}$
- (xii) If  $x$  is so small that its square and higher powers can be neglected, then show that  $\frac{1-x}{\sqrt{1+x}} \approx 1 - \frac{3}{2}x$



4.

Attempt any nine parts.

(2)  
MTN-42-21

9 × 2 = 18

- (i) If  $\sin \theta = -\frac{1}{\sqrt{2}}$   $R(\theta)$  is in 3<sup>rd</sup> quadrant. Find the value of  $\cot \theta$
- (ii) Verify that  $2\sin 45^\circ + \frac{1}{2} \operatorname{cosec} 45^\circ = \frac{3}{\sqrt{2}}$
- (iii) Verify that  $(\sec \theta + \tan \theta)(\sec \theta - \tan \theta) = 1$
- (iv) Express  $\sin 319^\circ$  as a trigonometric function of an angle of positive degree measure of less than  $45^\circ$ .
- (v) Prove that  $\tan(45^\circ + A) \cdot \tan(45^\circ - A) = 1$
- (vi) Prove that  $1 + \tan \alpha \cdot \tan 2\alpha = \sec 2\alpha$
- (vii) Find the period of  $3\cos \frac{x}{5}$
- (viii) Solve for  $C$  in a triangle  $\triangle ABC$  if  $\gamma = 90^\circ$ ,  $\alpha = 62^\circ 40'$  and  $b = 796$
- (ix) In an equilateral triangle find the value of  $R$ .
- (x) Prove that  $(r_1 + r_2) \tan \frac{\gamma}{2} = c$
- (xi) Find the value of  $\operatorname{cosec}(\tan^{-1}(-1))$
- (xii) Solve  $\sin x + \cos x = 0$  for  $x \in [0, 2\pi]$
- (xiii) Find the solution of  $\cot \theta = \frac{1}{\sqrt{3}}$  for  $\theta \in [0, \pi]$

**SECTION-II**

3 × 10 = 30

**NOTE: Attempt any three questions.**

- 5.(a) Solve the system of linear equations by Cramer's rule.

$$2x_1 - x_2 + x_3 = 8, \quad x_1 + 2x_2 + 2x_3 = 6, \quad x_1 - 2x_2 - x_3 = 1$$

- (b) If the roots of  $px^2 + qx + r = 0$  are  $\alpha$  and  $\beta$ , then prove that  $\sqrt{\frac{\alpha}{\beta}} + \sqrt{\frac{\beta}{\alpha}} + \sqrt{\frac{q}{p}} = 0$

- 6.(a) Resolve
- $\frac{3x+7}{(x^2+4)(x+3)}$
- into partial fractions.

- (b) Sum of three numbers in A.P. is 24 and their product is 440. Find the numbers.

- 7.(a) If
- $y = \frac{1}{3} + \frac{1 \cdot 3}{2!} \left(\frac{1}{3}\right)^2 + \frac{1 \cdot 3 \cdot 5}{3!} \left(\frac{1}{3}\right)^3 + \dots$
- then prove that
- $y^2 + 2y - 2 = 0$

- (b) Find the values of  $n$  and  $r$  when  ${}^nC_r = 35$  and  ${}^nP_r = 210$

- 8.(a) Find the values of the trigonometric function
- $\frac{-17\pi}{3}$

- (b) Prove that  $\frac{2\sin \theta \sin 2\theta}{\cos \theta + \cos 3\theta} = \tan 2\theta \tan \theta$

- 9.(a) Solve the triangle
- $ABC$
- if
- $a = 53$
- ;
- $\beta = 88^\circ 36'$
- ;
- $\gamma = 31^\circ 54'$

- (b) Prove that  $\sin^{-1}\left(\frac{5}{13}\right) + \sin^{-1}\left(\frac{7}{25}\right) = \cos^{-1}\left(\frac{253}{325}\right)$