

| Paper Code Number: 2183 | | 2023 (1 st -A) INTERMEDIATE PART-1 (11 th Class) | | Roll No: _____ | |
|-------------------------------------|--|---|------------------------------|--|------------------------------|
| STATISTICS PAPER-I MTN-11-23 | | | | | |
| TIME ALLOWED: 20 Minutes | | OBJECTIVE | | MAXIMUM MARKS: 17 | |
| Q.No.1 | You have four choices for each objective type question as A, B, C and D. The choice which you think is correct, fill that bubble in front of that question number, on bubble sheet. Use marker or pen to fill the bubbles. Cutting or filling two or more bubbles will result in zero mark in that question. | | | | |
| S.# | QUESTIONS | A | B | C | D |
| 1 | In binomial distribution $n = 16$ and $p = \frac{1}{2}$, then variance will be: | 2 | 4 | 5 | 6 |
| 2 | The hypergeometric experiment has properties: | One | Two | Three | Four |
| 3 | In plural sense, statistics mean: | Methods | Sample values | Numerical data | Average values |
| 4 | A pie chart is represented by: | Square | Circle | Triangle | Rectangle |
| 5 | The G.M. of 1, 3, and 27 is: | 10 | 27 | 10.3 | 3 |
| 6 | If $\bar{X} = 10$ and $Y = 2X + 7$, then $\bar{Y} = ?$ | 27 | 37 | 20 | 17 |
| 7 | For a set of 20 values, $\sum (X - \bar{X})^2 = 780$, then S.D. will be: | 49 | 7 | 14 | 98 |
| 8 | $Var(aX + b)$ equals to: | $Var(X) + b$ | $a Var(X) + b$ | $a^2 Var(X)$ | $Var(X)$ |
| 9 | The median of data -2, 0, 2, 5, -1 is: | -2 | 2 | 5 | 0 |
| 10 | For a normal distribution, $\bar{X} \pm 3S$ include of the observations: | 99.73% | 95.45% | 88.27% | 68.27% |
| 11 | Simple aggregate index number is given by: | $\frac{\sum P_0}{\sum P_n} \times 100$ | $\frac{P_n}{P_0} \times 100$ | $\frac{\sum P_n}{\sum P_0} \times 100$ | $\frac{P_0}{P_n} \times 100$ |
| 12 | Simple index number involves commodity: | Four | Two | Three | One |
| 13 | The probability of a black queen from a pack of 52 playing cards is: | $\frac{4}{52}$ | $\frac{2}{52}$ | $\frac{1}{52}$ | $\frac{3}{52}$ |
| 14 | If $P(A) = 0.4$, $P(B) = 0.5$, $P(A \cap B) = 0.2$ then $P(A \cup B) = ?$ | 0.7 | 0.8 | 0.6 | 0.5 |
| 15 | For a discrete random variable X , $\sum P(x)$ is always equal to: | 0 | 1 | 2 | 3 |
| 16 | If $Var(X) = 10$ and $Var(Y) = 20$, then $Var(X - Y) = ?$ | -10 | 20 | 10 | 30 |
| 17 | The hypergeometric distribution has parameters: | One | Two | Three | Four |

STATISTICS PAPER-I

TIME ALLOWED: 2.40 Hours

SUBJECTIVE

MAXIMUM MARKS: 68

NOTE: Write same question number and its parts number on answer book, as given in the question paper.

SECTION-I

MTN-11-23

2. Attempt any eight parts.

8 × 2 = 16

- (i) Distinguish between the terms population and sample. (ii) Narrate any two sources of collecting primary data.
- (iii) Explain the term weighted mean with formula. (iv) Write down the names of any four positional averages.
- (v) Describe the empirical relation between mean, median and mode, for moderately skewed distribution.
- (vi) Given that $X_1 = 3$ and $X_2 = 27$. Show that G.M > H.M.
- (vii) Find the modal letter of the word "PAKISTAN".
- (viii) Given that $U = \frac{X-98}{5}$, $\sum fU = -30$ and $\sum f = 30$. Find \bar{X} .
- (ix) Explain the concept of unweighted index number.
- (x) Define Laspeyre's price index number with formula.
- (xi) Given that $\sum p_1q_1 = 1400$, $\sum p_2q_2 = 1600$, $\sum p_0q_1 = 1360$ and $\sum p_0q_2$. Compute Paasche's price index numbers.
- (xii) If link relatives are 100, 102, 113 and 118. Find chain indices.

3. Attempt any eight parts.

8 × 2 = 16

- (i) What is main idea of classification? (ii) Distinguish between class limits and class boundaries.
- (iii) Enlist the types of dispersion. (iv) Outline any two properties of S.D.
- (v) How would you explain the concept of Kurtosis if $b_2 > 3$, $b_2 = 3$ and $b_2 < 3$?
- (vi) Find b_1 , if $m_2 = 5.2$ and $m_3 = -0.8$ (vii) The first two moments about 4 are 1 and 16. Find variance.
- (viii) Is variance affected by change of origin and scale?
- (ix) Describe the main idea of calculating probability of an event.
- (x) Distinguish between sample space and sample point. (xi) What is the range of probability of an event?
- (xii) Two cards are drawn without replacement from 52 playing cards. What is the probability that both are aces?

4. Attempt any six parts.

6 × 2 = 12

- (i) Define probability distribution. (ii) Explain the application of random numbers.
- (iii) Given $f(x) = \frac{k}{x}$, $x = 1, 2, 3$. Find k . (iv) Given $E(X) = 200$ and $S.D(X) = 5$. Find $E(X^2)$
- (v) What are the parameters of a binomial distribution?
- (vi) A random variable X has following binomial distribution with $n = 5$ and $p = 0.2$. Find $P(X = 2)$
- (vii) In a binomial distribution, mean = 2.4 and standard deviation = 1.2. Find the value of n .
- (viii) Define hypergeometric experiment.
- (ix) For hypergeometric distribution $N = 10$, $n = 4$ and $K = 5$. Find $P(X = 0)$

SECTION-II

NOTE: Attempt any three questions.

3 × 8 = 24

- 5.(a) The average wage of 4 men is Rs.17 per hour. What is the average wage of further 6 men if the average wage of all 10 men is Rs.20 per hour. 4
- (b) Compute G.M of the data 4
- | | | | | | | |
|-----|-------|-------|-------|-------|-------|-------|
| Age | 11-20 | 21-30 | 31-40 | 41-50 | 51-60 | 61-70 |
| f | 12 | 14 | 26 | 35 | 23 | 5 |
- 6.(a) Find semi-inter quartile range for the data given below: 4
- | | | | | | |
|------|----|----|-----|----|----|
| Ages | 20 | 30 | 40 | 50 | 60 |
| f | 3 | 61 | 132 | 51 | 2 |
- (b) Given the first four moments about $Y = 20$ are as $-2, 15, -25$ and 80 respectively. Calculate b_2 . 4
- 7.(a) Compute: (i) Base year Weighted Index Number (ii) Current year Weighted Index Number from the given data by taking 1980 as base year. 4
- | Commodities | Prices | | Quantities | |
|-------------|--------|------|------------|------|
| | 1980 | 1981 | 1980 | 1981 |
| A | 10 | 12 | 20 | 22 |
| B | 8 | 8 | 16 | 18 |
| C | 5 | 6 | 10 | 11 |
| D | 4 | 5 | 7 | 8 |
- (b) A bag consists of 3 white and 5 red balls. If two balls are drawn at random, what is the probability that: 4
- (i) Both are white (ii) Both are of same colour
- 8.(a) A random variable ' X ' has the following probabilities distribution: 4
- | | | | |
|--------|---------------|---------------|---------------|
| X | 1 | 2 | 3 |
| $P(x)$ | $\frac{6}{9}$ | $\frac{2}{9}$ | $\frac{1}{9}$ |
- Find Mean and Standard Deviation of x .
- (b) A continuous random variable ' X ' can assume values between $X = 2$ and $X = 5$ and has a density function $f_x = \frac{2(1+X)}{27}$ (i) $P(3 < x < 4)$ (ii) $P(X \leq 4)$ 4
- 9.(a) Let ' X ' be a random variable having binomial distribution with parameters $n = 5$ and $p = 0.52$ Calculate (i) $P(X = -2)$ (ii) $P(X = 2)$ (iii) $P(X = 2.5)$ (iv) $P(X > 5)$ 4
- (b) A committee of size "3" is to be selected at random from 4 women and 6 men. Obtain the probability distribution of number of men in the committee. 4