

☆☆☆☆

Inter - (Part-I) - A-2022

Roll No. _____ to be filled in by the candidate

(For All Sessions)

Paper Code	6	1	8	8
------------	---	---	---	---

Statistics (Objective Type)

Time: 20 Minutes

Rwp 22

Marks: 17

NOTE: Write answers to the questions on objective answer sheet provided. Four possible answers A, B, C & D to each question are given. Which answer you consider correct, fill the corresponding circle A, B, C or D given in front of each question with marker or pen ink on the answer sheet provided.

- 1.1. The probability of sure event is.

(A) 0	(B) 1	(C) >1	(D) <1
-------	-------	----------	----------
2. The amount of milk produced by cow is _____ variable.

(A) Discrete	(B) Continuous	(C) Qualitative	(D) None
--------------	----------------	-----------------	----------
3. If $E(X)=4$, the arithmetic Mean will be.

(A) 4	(B) Zero	(C) 8	(D) 1
-------	----------	-------	-------
4. In binomial experiments, each trial has:

(A) One outcome	(B) Two outcomes	(C) Three outcomes	(D) Four outcomes
-----------------	------------------	--------------------	-------------------
5. In hypergeometric distribution $N = 6, n = 2, K = 3$, then mean is:

(A) 2	(B) 3	(C) 1	(D) 4
-------	-------	-------	-------
6. The grouped data are also called.

(A) Raw data	(B) Primary data	(C) Secondary data	(D) Qualitative data
--------------	------------------	--------------------	----------------------
7. The average value of a lower and upper limits of a class is called:

(A) Class boundary	(B) Class frequency	(C) Mid point	(D) Class interval
--------------------	---------------------	---------------	--------------------
8. Graph of time series is known as:

(A) Histogram	(B) Ogive	(C) Historigram	(D) Polygon
---------------	-----------	-----------------	-------------
9. Geometric Mean of the values 2, 4, -3, 6, 0 is:

(A) -3	(B) 0	(C) 3	(D) Cannot be computed
--------	-------	-------	------------------------
10. We must arrange the data before calculating:

(A) Mode	(B) Median	(C) Mean	(D) G.M
----------	------------	----------	---------
11. If 10% is added to each value of variable, the geometric mean of new variable is added by:

(A) 10%	(B) No change	(C) 10	(D) 110
---------	---------------	--------	---------
12. Variance remains unchanged by change of:

(A) Scale	(B) Origin	(C) Both (A) and (B)	(D) None
-----------	------------	----------------------	----------
13. A measure of dispersion is always:

(A) Zero	(B) Positive	(C) Negative	(D) None of these
----------	--------------	--------------	-------------------
14. Second moment about Mean is called:

(A) Mean	(B) S.D	(C) C.V.	(D) Variance
----------	---------	----------	--------------
15. In chain base method, the base period is:

(A) Fixed	(B) Changed	(C) Constant	(D) None of these
-----------	-------------	--------------	-------------------
16. Base year weighted index numbers are:

(A) Laspeyre's Index	(B) Paasche's Index	(C) Fisher Index	(D) Marshall Index
----------------------	---------------------	------------------	--------------------
17. The probability of an event always lies between:

(A) Zero and 2	(B) -1 and +1	(C) Zero and 1	(D) -2 and +2
----------------	---------------	----------------	---------------

R

Roll No. _____ to be filled in by the candidate

(For All Sessions)

Statistics (Essay Type)

Time: 2:40 Hours

Section - I

Marks:68

2 x 8 = 16

2- Write short answers of any eight parts from the following.

- Define statistics.
- Find the G.M from the following values. 4, 5, 10, 0, 20.
- Write down the advantages of mode.
- If sum of deviation from $X = 15$ for 10 values is 25, then find A.M.
- Define composite index number.
- If $\sum P_o q_o = 362, \sum P_1 q_o = 428, \sum P_o q_1 = 398, \sum P_1 q_1 = 470$ then find Fisher's Ideal I.No.

3- Write short answers of any eight parts from the following.

- Define "Histogram".
 - Define Quartile deviation.
 - Define range. Also give an example.
 - Compute coefficient of standard deviation if Mean = 125 and standard deviation = 2.
 - Make a sample space if we toss a fair coin three times.
 - Give the statement of addition Law of probability for two non-mutually exclusive events.
- 4- Write short answers of any six parts from the following.
- What are random numbers?
 - What is probability density function?
 - If $E(X) = 3$ and $E(Y) = 2.5$, then find $E(X+Y)$.
 - What are parameters of binomial distribution?
 - In hypergeometric distribution $n = 5, K = 4$ and $N = 12$ then find its mean.

- Distinguish between discrete variable and continuous variable.
- Define Median.
- What are merits of mode?
- Define weighted mean.
- If paasche's I.No = 74.76 and Fishers I.No = 75.76 then find Laspayer's I.No = ?
- Define link relative.

2 x 8 = 16

- Define relative frequency.
- Compute coefficient of quartile deviation, if $Q_1 = 12.50$ and $Q_3 = 48.36$
- Define mean deviation.
- Compute mean coefficient of dispersion if mean deviation = 3.92 and Mean = 16.25
- How many permutations can be formed from the word "STATISTICS"?
- State the multiplication law of probability for independent events.

2 x 6 = 12

Section - II

8 x 3 = 24

NOTE : Answer any three questions from the following.

- 5.(a) The frequency distribution given below has $D = X - 8$ Find the Geometric Mean.

D	-12	-8	-4	0	4	8	12	16
f	2	5	8	18	22	13	8	4

- 6.(a) Compute mean deviation from the data given below using mean.

Classes	5-9	10-14	15-19	20-24	25-29
f	5	8	12	10	5

- 7.(a) Compute Fisher's index number for the following data.

commodities	Base year		Current year	
	Price	Quantity	Price	Quantity
A	7	70	5	49
B	5	27	7	28
C	10	35	9	29
D	9	50	4	42

04+04

- (b) Compute the median and mode of the following distribution.

Classes	0-7	7-14	14-21	21-28	28-35
f	5	8	7	15	5

- (b) Calculate Bowley's coefficient of skewness for the following data.

Groups	2-4	4-6	6-8	8-10	10-12
frequency	3	5	7	3	2

04+04

- (b) A pair of dice is thrown. Find the probability of getting a total of either 5 or 11.
- 8.(a) Find the missing value of 'A' from the following probability distribution.

x	2	3	4	5	6
P(x)	0.01	0.25	0.40	A	0.04

Also find $E(x)$

- 9.(a) A fair coin is tossed 5 times. What is the probability of getting.
- Exactly 3 heads
 - At least 3 heads

- (b) A continuous random variable X has a density function. $f(x) = 2x, 0 \leq x \leq 1$ find $P(0 < x < 0.2)$

04+04

- (b) Find $P(x \leq 2)$ for hypergeometric distribution having $N = 8, K = 5, n = 6$

04+04